

## 1. Selecting a site location

Consider the following requirements when selecting a site location for your offsite-indoor flu clinic.

<b>Clinic Flow</b>	<ul style="list-style-type: none"> <li>Space to enable one-way foot traffic:               <ul style="list-style-type: none"> <li>Separate entry and exit points</li> </ul> </li> <li>Space for clinic functions:               <ul style="list-style-type: none"> <li>Screening and registration</li> <li>Vaccine storage and preparation</li> <li>Vaccination administration</li> <li>Waiting areas for post-vaccination observation</li> <li>Emergency care</li> </ul> </li> <li>Space for parking</li> <li>Space for traffic flow into/out of location</li> </ul>
<b>Utilities</b>	<ul style="list-style-type: none"> <li>Adequate heating and cooling</li> <li>Adequate lighting</li> <li>Internet access</li> <li>Power outlets (for portable vaccine refrigerators, etc.)</li> </ul>
<b>Accessibility</b>	<p>Public spaces must comply with the <a href="#">AODA standards for Accessible Public Spaces</a>.</p> <ul style="list-style-type: none"> <li>Accessible paths and walkways</li> <li>Accessible entry/exit</li> <li>Accessible restrooms</li> <li>Accessible waiting areas</li> <li>Proximity to population centers and mass transit</li> </ul>

## 2. Setting up the clinic

<b>Flow and layout</b>	<ul style="list-style-type: none"> <li>Designate and set up spaces in clinic:               <ul style="list-style-type: none"> <li>Separate entrance and exit</li> <li>Screening/temperature check station at entrance</li> <li>Waiting area (chairs at least 6 feet apart)</li> <li>Vaccine preparation area (separate from vaccination stations)</li> <li>Vaccination stations (at least 6 feet apart, designated stations for each vaccine type offered, if applicable)</li> <li>Area for special-needs patients (e.g., persons with disabilities or limited mobility)</li> <li>Area for evaluation/treatment of patients with adverse events or other medical needs</li> <li>Area for staff to leave personal items and take breaks</li> <li>Area for administrative work and documentation</li> </ul> </li> <li>Put up signage:               <ul style="list-style-type: none"> <li>Screening signage at entrance</li> <li>Signs, banners, and floor markers to guide traffic in one direction</li> <li>Signs with public health messaging (physical distancing, hand hygiene, respiratory hygiene, cough etiquette)</li> </ul> </li> </ul>
<b>Data management</b>	<p>Develop a data collection and management strategy based on site capability (manual processes must be planned for temporary sites lacking specific infrastructure). Consider:</p> <ul style="list-style-type: none"> <li>Access to vaccination history (if applicable)</li> <li>Reporting to EMR</li> <li>Reporting to patient's primary care provider</li> </ul>

## 2. Setting up the clinic (continued)

<b>Supplies</b>	<ul style="list-style-type: none"> <li>Alcohol-based hand sanitizer (at least 70% alcohol)</li> <li>Hand soap</li> <li>Cleaning supplies</li> <li>PPE for staff (masks, eye protection, gloves)</li> <li>Masks for patients who come without</li> <li>Thermometers (if performing temperature checks at screening station)</li> <li>Tissues</li> <li>Adhesive bandages or tape/cotton balls for injection site</li> <li>Screening and documentation forms</li> <li>Vaccines and diluents (if needed)</li> <li>Prep pads</li> <li>Sterile alcohol wipes (individually packaged)</li> <li>Syringes</li> <li>Needles (varying lengths appropriate for the expected patient population)</li> </ul>	<ul style="list-style-type: none"> <li>Sharps containers (closable, puncture-resistant, and leakproof)</li> <li>Emergency medical kit with epinephrine with signed medical orders</li> <li>First aid kit</li> <li>Disposable table covers that can be changed if soiled</li> <li>Technology (if using):               <ul style="list-style-type: none"> <li>Computers / tablets</li> <li>Printers</li> <li>2D barcode readers</li> <li>Internet access or hotspots</li> </ul> </li> <li>Outlet strips (multi-plug) and extension cords</li> <li>Office supplies (including pens, printer paper, etc.)</li> <li>Wastebaskets</li> </ul>
-----------------	--	---

## 3. Operating the clinic

<b>Scheduling and booking appointments</b>	<ul style="list-style-type: none"> <li>Schedule multiple smaller clinics instead of large clinics to avoid crowds</li> <li>Schedule appointments (rather than allowing walk-ins)</li> <li>Request that only one adult accompanies children for their appointments</li> <li>Have patients wait outside until their appointment time (if appropriate) and call them when ready</li> </ul>
<b>Staffing</b>	<ul style="list-style-type: none"> <li>Schedule extra clinic staff and volunteers for:               <ul style="list-style-type: none"> <li>Monitoring traffic flow and waiting areas</li> <li>Screening</li> <li>Registration and consent processes</li> <li>Cleaning</li> </ul> </li> </ul>
<b>Cleaning</b>	<ul style="list-style-type: none"> <li>Clean surfaces and equipment (including the rolling carts) between patients, and at the beginning and end of every shift. Follow COVID-19 environmental cleaning protocols for clinic settings (<a href="#">BCCDC, 2020</a>)</li> </ul>
<b>Screening</b>	<ul style="list-style-type: none"> <li>Implement passive screening procedures at entry with signage</li> <li>Implement active screening procedures at entry. See Ontario COVID-19 Screening Document for screening checklist. (<a href="#">MOH, June 11, 2020</a>)</li> <li>Divert screen-positive patients to seek testing at an assessment centre or emergency room as appropriate:               <ul style="list-style-type: none"> <li>Ask the patient to perform hand hygiene</li> <li>Provide the patient with a medical mask</li> <li>Make efforts to ensure the patient has a method of travel that maintains physical distancing</li> </ul> </li> </ul>
<b>Contactless communications</b>	<ul style="list-style-type: none"> <li>Use an online or telephone system to:               <ul style="list-style-type: none"> <li>make appointments</li> <li>collect registration information</li> <li>screen patients</li> <li>provide vaccine information in advance</li> </ul> </li> </ul>

**3. Operating the clinic (continued)**

**Contactless communications**

Complete information forms on behalf of patients. If signature is required:  
 Provide a separate pen for each patient  
 Clean pens between each use  
 Provide pre-immunization information on posters, videos, and/or QR codes. Ensure that information is accessible (available in multiple languages as needed)

**4. Maintaining the cold chain**

Adapted from Vaccine Storage and Handling Guidelines ([Government of Ontario, 2018](#))

The “cold chain” includes all materials, equipment and procedures used to maintain vaccines in the required temperature range of +2°C to +8°C from the time of manufacture until the vaccines are administered to individuals. Most vaccines are considered to be damaged at 0°C. The loss of vaccine effectiveness due to exposures to adverse conditions is cumulative, permanent and irreversible.

**Setup**

**Staff**

Designate one person as the **lead** to monitor vaccine storage and handling practices.  
 Designate one person as **back-up** to monitor vaccine storage and handling practices.  
 Ensure designated staff understand rationale for correct vaccine storage and handling.

**Equipment**

Temperature Log Book to track:  
 Day  
 Time of each temperature check  
 Current temperature  
 Maximum temperature  
 Minimum temperature  
 Name of staff member performing temperature check

Refrigerator:  
 Purpose-built refrigerators are recommended for storing large inventories of vaccines. They ensure maintenance of +2°C to +8°C internal temperature; ongoing air circulation ensures; evaporator to prevent vaccine from freezing; temperature recovery system.  
 Domestic or bar refrigerators are not recommended for vaccine storage. It is complicated to manage their temperatures, which makes them unsuitable for vaccine storage unless excellent vaccine storage and handling practices are followed.  
 If using a domestic/bar fridge: contact your public health unit for assistance on required modifications prior to vaccine storage in order to ensure that vaccines will be safely stored.  
 Temperature monitoring device for each vaccine refrigerator and insulated container.  
 Insulated containers for temporary storage in case of power outage.

**Refrigerator setup**

Stabilize the temperature of the vaccine refrigerator before stocking vaccine.  
 Place the maximum-minimum thermometer probe on the middle refrigerator shelf inside an empty vaccine box.  
 Place the refrigerator out of direct sunlight.  
 Mark the plug clearly so the refrigerator is not unplugged or turned off accidentally.  
 Keep icepacks/gel packs in your freezer

**4. Maintaining the cold chain (continued)**

**Setup**

**Receiving and returning vaccines**

When you receive your order, check that you received your full order and ensure that the order matches the packing slip.  
 Place vaccines in the refrigerator immediately.  
 Store vaccines in their original packaging.  
 DO NOT store vaccines in the door of the refrigerator.  
 If you did not receive everything you ordered, or your order does not match the packing slip, notify your vaccine supply source (i.e., public health units or OGPMS) immediately.  
 Always return expired or spoiled vaccine to your vaccine supply source for disposal. The Ministry may be reimbursed by manufacturers for returned vaccines.

**Cold chain management**

**Daily tasks**

Ensure your refrigerator is in good working order.  
 Aim to maintain vaccine refrigerator temperature at +5°C, as this gives a greater leeway for protection from temperature fluctuation.  
 Keep your refrigerator door openings to a minimum.  
 Check temperatures. Temperature monitoring and recording devices do not guarantee safety of vaccines and are not to be considered a substitute for the manual recording of minimum, maximum and current temperatures twice daily.  
     If using refrigerators, check and record the maximum, minimum and current temperatures twice daily in the Temperature Log Book at the beginning and end of each day.  
     If using insulated containers as temporary storage, see "Management of insulated containers" below for temperature check schedule.  
 Reset maximum-minimum thermometer after recording the temperature readings.  
 When using a data logger, the minimum, maximum and current temperatures still need to be recorded manually as a timely alert to any breach in the cold chain.  
 Do not store food, beverages or medical/laboratory specimens in the vaccine refrigerator.  
 Fill the lower drawers and the door with water bottles.  
 Do not overstock your refrigerator with vaccines.  
 Rotate stock so vaccines with the shortest expiry date are used first.  
 If vaccines have been exposed to temperatures below +2°C or above +8°C, contact your public health unit immediately.

**Power outage**

Always have an alternative means of vaccine storage available.  
 In the Temperature Log Book, document the time and the maximum, minimum and current temperature inside of the non-functioning refrigerator.  
 Do not allow the vaccine to remain in a non-functioning unit for an extended period of time.  
 Record the time and refrigerator temperature when the electricity supply is restored, and again when the thermometer reading is within +2°C to +8°C.  
 Call your public health unit immediately to report any exposures to temperatures below +2°C and/or above +8°C.  
 Segregate the exposed vaccines in the refrigerator by placing these vaccines in a labelled container (or bag), marked with the date and time and "DO NOT USE."  
 Never use or discard the vaccine until your public health unit has assessed the situation.

## Temporary storage: insulated containers

Properly packed insulated containers (coolers) can maintain the required temperature for 3-4 hours, subject to environmental and physical conditions. Use insulated containers for:

- Transporting vaccine
- Temporary storage of vaccine (during clinics or when cleaning the refrigerator)
- Emergency storage (power outage)

### Management of insulated containers

Experiment to find the correct combination of icepack(s) and/or gel pack(s) to ensure each insulated container is able to maintain the required temperatures. See below for instructions on how to properly pack an insulated container.

Do not place insulated containers with vaccines in the trunk of a car.

Visually inspect the temperature every time the insulated container is opened.

Record temperature in the insulated container:

- Before leaving office
- After 1 hour of travel
- Upon arrival at clinic
- Every hour during the session
- Upon completion of the session at the clinic
- Upon arrival back to office

## Packing insulated containers

• Ice/gel packs must be correctly conditioned before use. See Preparation instructions below. The risk of freezing vaccines increases if the icepacks/gel packs are not correctly conditioned.

- Incorrect use of gel packs is riskier than icepacks because gel packs remain colder than 0°C for longer.
- Freezing episodes happen very easily in all coolers, usually in the first 2 hours after packing.

Item	Preparation	Use Instructions
<b>Insulated hard sided container</b>	Pre-chill insulated container until a temperature between +2°C to +8°C is reached prior to placing vaccines into the container. Use frozen gel packs or place the container in a refrigerator.	Check container for damage. Ensure container can close fully and maintain its seal.
<b>Vaccine and temperature monitoring device</b>	Check the accuracy of devices once annually and change the battery every 6 months.	Position maximum-minimum thermometer sensor inside vaccine box.
<b>Inner flexible ice blanket</b>	Condition in refrigerator at +2°C to +8°C	Wrap inner flexible ice blanket around vaccines.
<b>Gel packs</b>	Condition gel packs in freezer to appropriate temperature for season: <ul style="list-style-type: none"> <li>• Winter transport: +2°C to +8°C.</li> <li>• Summer transport: -10°C to -20°C.</li> </ul>	Place gel packs on top of outer flexible ice blanket. Additional icepacks may be required depending on cold-life needed for the length of transport.
<b>Outer flexible ice blanket</b>	Condition in refrigerator at +2°C to +8°C	Wrap outer flexible ice blanket around vaccines and inner flexible ice blanket.
<b>Additional insulating material</b>	N/A	Bubble wrap, Styrofoam chips, crumpled or shredded newspaper should be placed inside (bottom, top and sides) the insulated container to allow for cool air circulation.

*Offsite-Indoor Clinic Checklist is adapted from: [Guidance on seasonal influenza and antiviral drug use in Canada for the 2020-2021 influenza season in the context of COVID-19 \(NCCID, Sept 2020\)](#); [Guidance on the use of influenza vaccine in the presence of COVID-19 \(NACI, Sept 29, 2020\)](#); [Guidance for influenza vaccine delivery in the presence of COVID-19 \(NACI, August 5, 2020\)](#); [Considerations for Planning Curbside/Drive-Through Vaccination Clinics \(CDC, July 27, 2020\)](#)*



**Offsite-Indoor Clinic Checklist** is a product of the Centre for Effective Practice. Permission to use, copy, and distribute this material for all non-commercial and research purposes is granted, provided the above disclaimer, this paragraph and the following paragraphs, and appropriate citations appear in all copies, modifications, and distributions. Use of the Offsite-Indoor Clinic Checklist for commercial purposes or any modifications of the Resource are subject to charge and use must be negotiated with the Centre for Effective Practice (Email: [info@cep.health](mailto:info@cep.health))

For statistical and bibliographic purposes, please notify the Centre for Effective Practice ([info@cep.health](mailto:info@cep.health)) of any use or reprinting of the Resource. Please use the below citation when referencing the Resource:

Reprinted with Permission from the Centre for Effective Practice. (Oct 2020). Offsite-Indoor Clinic Checklist: Ontario. Toronto: Centre for Effective Practice.

Developed by:



Centre  
for Effective  
Practice

With input and support from:



This resource was developed with input from and support by Ontario College of Family Physicians (OCFP), Association of Family Health Teams of Ontario (AFHTO), Ontario Medical Association (OMA) and OMA SGFP, Nurse Practitioners' Association of Ontario (NPAO) and Registered Nurses' Association of Ontario (RNAO).

It has been reviewed and endorsed by the Provincial Primary Care Advisory Table (PPCAT) established by the Ministry of Health.